

On simple ringoids

JENS ZUMBRÄGEL*

Abstract. A ringoid is a set with two binary operations that are linked by the distributive laws. We study special classes of ringoids that are congruence-simple or ideal-simple. In particular, we examine generalised parasemifields and non-associative semirings.

Keywords: ringoid, semiring, parasemifield, congruence-simple, ideal-simple

2000 Mathematics Subject Classification: 16Y99

1. Introduction

A *ringoid* $(S, +, *)$ is defined as a set S with two binary operations, $+$ and $*$, such that $a*(b+c) = a*b + a*c$ and $(a+b)*c = a*c + b*c$ for all $a, b, c \in S$. In other words, a ringoid is the most general ring-like structure in which the distributive laws hold.

In this paper a ringoid $(S, +, *)$ is called a *semiring* if the groupoid $(S, +)$ is a commutative semigroup; if $(S, +)$ is even an abelian group we speak of a *ring*. Semirings or rings $(S, +, *)$ in which the groupoid $(S, *)$ is a semigroup are called *associative*. If we have a ringoid $(S, +, *)$ in which $(S, *)$ is a quasigroup we speak of a *generalised parasemifield*; if in addition $(S, +, *)$ is a semiring the ringoid $(S, +, *)$ is called a *parasemifield*.

In ringoids the notions of ideal and congruence can be defined (see Section 2) and we can speak of *ideal-simple* and *congruence-simple* ringoids. In the last years there has been a considerable interest in congruence-simple associative semirings. The classification in the commutative case was given in [EHJK01], and finite non-commutative semirings with zero were classified in [Zum08]. Studies of general congruence-simple associative semirings can be found e.g. in [Mon04, EK07]. On the other side, parasemifields appear to be an important class of ideal-simple semirings and a closer investigation of commutative parasemifields has been started recently [KK08, KKK08].

The aim of the present paper is to initiate the study of more general ringoids that are congruence-simple or ideal-simple. We will be mainly interested in ringoids in which at least one of the binary operation is non-associative.

The outline of this paper is as follows. In Section 2 we give the necessary definitions and present some preliminary results. We present a basic classification of ideal-simple ringoids, provided that the multiplication is associative and

*School of Mathematical Sciences, University College Dublin, Belfield, Dublin 4, Ireland.
E-mail: jens.zumbragel@ucd.ie

This work has been supported by the Science Foundation Ireland under grant no. 08/IN.1/I1950.

commutative. In Section 3 we examine generalised parasemifields, where in particular we study the additive groupoid and its automorphism group. Finally, in Section 4 we investigate congruence-simple ringoids, where we restrict ourselves to the case of (not-necessarily associative) semirings. We give a basic classification of congruence-simple semirings and a characterisation of finite so-called k -ideal-simple semirings having idempotent addition.

2. Prerequisites

Groupoids

Definition 2.1. A **groupoid** $(S, \circ)$ is a non-empty set S with a binary operation $\circ : S \times S \rightarrow S$.

We introduce some notation. Let $\text{Sym}(S)$ be the symmetric group on the set S , and let $T(S)$ be the monoid of all maps $S \rightarrow S$. The group of automorphisms of $(S, \circ)$ is denoted by $\text{Aut}(S, \circ)$, and the monoid of endomorphisms of $(S, \circ)$ is denoted by $\text{End}(S, \circ)$. Clearly, $\text{Aut}(S, \circ)$ is a subgroup of $\text{Sym}(S)$ and $\text{End}(S, \circ)$ is a submonoid of $T(X)$.

For $a \in S$ consider the left and right multiplication maps L_a and R_a , defined by $L_a(x) = a \circ x$ and $R_a(x) = x \circ a$ for $x \in S$. The submonoid of $T(X)$ generated by $\{L_a \mid a \in S\} \cup \{R_a \mid a \in S\}$ is denoted by $\text{Mult}(S, \circ)$.

For subsets A and B of S let $A \circ B$ denote the subset $\{a \circ b \mid a \in A, b \in B\}$ of S . Furthermore, for $a, b \in S$ we will write $a \circ B$ and $A \circ b$ for the sets $\{a\} \circ B$ and $A \circ \{b\}$.

Definition 2.2. Let $(S, \circ)$ be a groupoid. A **congruence** of $(S, \circ)$ is an equivalence relation ρ on the set S such that $\rho(a, b)$ implies $\rho(x \circ a, x \circ b)$ and $\rho(a \circ x, b \circ x)$ for all $a, b, x \in S$.

A non-empty subset A of S is called a **subgroupoid** of the groupoid $(S, \circ)$ if $A \circ A \subseteq A$ holds. It is called an **ideal** of the groupoid $(S, \circ)$ if we have $(S \circ A) \cup (A \circ S) \subseteq A$.

A subgroupoid or ideal A of $(S, \circ)$ will be called *proper* if $A \neq S$.

Ringoids

Definition 2.3. A **ringoid** $(S, +, *)$ is a non-empty set S with two binary operations, $+$ and $*$, such that $a * (b + c) = a * b + a * c$ and $(a + b) * c = a * c + b * c$ hold for all $a, b, c \in S$.

As usual we call the operation $+$ *addition* and the operation $*$ *multiplication*.

Remark 2.4. Given two groupoids $(S, +)$ and $(S, *)$, then $(S, +, *)$ is a ringoid if and only if $\text{Mult}(S, *)$ is a submonoid of $\text{End}(S, +)$.

Simple ringoids

Definition 2.5. Let $(S, +, *)$ be a ringoid. A **congruence** on $(S, +, *)$ is an equivalence relation on the set S that is a congruence of both the groupoid $(S, +)$ and the groupoid $(S, *)$.

A non-empty subset A of S is called an **ideal** of the ringoid $(S, +, *)$ if A is a subgroupoid of the groupoid $(S, +)$ and an ideal of the groupoid $(S, *)$; that is, it holds $(A + A) \cup (S * A) \cup (A * S) \subseteq A$.

Definition 2.6. The ringoid $(S, +, *)$ is said to be **congruence-simple** if its only congruences are $\rho = \text{id}_S := \{(a, a) \mid a \in S\}$ and $\rho = S \times S$. It is said to be **ideal-simple** if there are no proper ideals A of $(S, +, *)$ with $|A| \geq 2$. If there are no proper ideals of $(S, +, *)$ at all, the ringoid will be called **ideal-free**.

We note that a ringoid $(S, +, *)$ is congruence-simple if and only if every non-constant homomorphism $(S, +, *) \rightarrow (T, +, *)$ into another ringoid $(T, +, *)$ is injective.

Ideal-simple ringoids

We present a basic classification of ideal-simple ringoids, provided that the multiplication is associative and commutative (note that these structures are formally akin to semirings, with the role of addition and multiplication interchanged).

We will use the following standard result:

Lemma 2.7. *Let $(S, *)$ be a semigroup such that $S * a = S = a * S$ for all $a \in S$. Then S is a group.*

Proposition 2.8. *Let $(S, +, *)$ be a ringoid with associative and commutative multiplication. If $(S, +, *)$ is ideal-simple then one of the following holds:*

- (i) $|S * S| = 1$,
- (ii) $(S, *)$ is a group,
- (iii) there is an absorbing element o in $(S, *)$ such that $(S \setminus \{o\}, *)$ is a group.

Proof. We may assume $|S| \geq 2$. Consider the set

$$N := \{a \in S \mid |S * a| = 1\},$$

which is either empty or an ideal of $(S, +, *)$. If N is empty then for all $a \in S$ we have $|S * a| > 1$, and, since $S * a$ is an ideal of $(S, +, *)$, we have $S * a = S$. Consequently, $(S, *)$ is a group. If $N = S$ we have $|S * a| = 1$ for all $a \in S$ and it follows that $|S * S| = 1$.

Suppose now that N is non-empty and $N \neq S$. Then $|N| = 1$, so $N = \{o\}$ for some $o \in S$, and we have $S * o = \{c\}$ for some $c \in S$. Furthermore, for all $a \neq o$

we have $S * a = S$. Let $T := S \setminus \{o\}$ and $a, b \in T$. Then $S = S * b = S * a * b$, so that $a * b \neq o$. Since $S * a = S$ and $S * o = \{c\}$ we have $S \setminus \{c\} \subseteq T * a \subseteq T$. This proves that $c = o$ and $T * a = T$ for all $a \in T$. Hence o is absorbing in $(S, *)$ and $(T, *)$ is a group. $\square$

In the next section we examine ringoids $(S, +, *)$ in which the multiplication is a group or, more generally, a quasigroup in more detail.

3. Generalised parasemifields

A groupoid $(S, *)$ is called a *quasigroup* if for all $a \in S$ the multiplication maps $L_a : x \mapsto a * x$ and $R_a : x \mapsto x * a$ are bijective maps of S , i.e. $L_a, R_a \in \text{Sym}(S)$. In this case $\text{Mult}(S, *)$ is a submonoid of $\text{Sym}(S)$.

Definition 3.1. A *generalised parasemifield* is a ringoid $(S, +, *)$ such that $(S, *)$ is a quasigroup.

Apparently, every generalised parasemifield is an ideal-free ringoid.

Remark 3.2. Given two groupoids $(S, +)$ and $(S, *)$, then $(S, +, *)$ is a generalised parasemifield if and only if $\text{Mult}(S, *)$ is a submonoid of $\text{Aut}(S, +)$.

The additive automorphism group

Lemma 3.3. Let $(S, +, *)$ be a generalised parasemifield. Then the additive automorphism group $\text{Aut}(S, +)$ is transitive.

Proof. Let $x, y \in S$ be given. Since $R_x \in \text{Sym}(S)$ there exists $a \in S$ such that $R_x(a) = y$. Now $L_a \in \text{Aut}(S, +)$ and we have $L_a(x) = a * x = R_x(a) = y$. $\square$

Corollary 3.4. Let $(S, +, *)$ be a generalised parasemifield, $|S| > 1$. Then $(S, +)$ has neither a neutral element nor an absorbing element.

Proof. If $(S, +)$ had a neutral resp. absorbing element then, since $\text{Aut}(S, +)$ is transitive, every element of S would be neutral resp. absorbing in $(S, +)$. But each groupoid can have at most one neutral or absorbing element. $\square$

Corollary 3.5. Let $(S, +, *)$ be a generalised parasemifield such that $(S, +)$ is a commutative semigroup. Then $|S| = \infty$.

Proof. Suppose $|S| < \infty$. It is well-known that every finite semigroup has an idempotent element, hence $(S, +)$ has an idempotent element. Since $\text{Aut}(S, +)$ is transitive, every element of S is idempotent. It follows that $(S, +)$ has an absorbing element, namely $\sum_{x \in S} x$, in contradiction to Corollary 3.4. $\square$

Example 3.6. The ringoid $(\mathbb{Z}, \max, +)$ is an example of an infinite generalised parasemifield with associative and commutative addition.

Groupoids with full automorphism group

The discussion of the automorphism group of groupoids is motivated by Remark 3.2 and Lemma 3.3.

If for a groupoid $(S, +)$ every bijection of S is an automorphism, i.e. $\text{Aut}(S, +) = \text{Sym}(S)$, then for any quasigroup $(S, *)$ we get a generalised parasemifield $(S, +, *)$. A proof of the following result can be found in [Tam63, Tam67].

Proposition 3.7. *A groupoid $(S, \circ)$ satisfies $\text{Aut}(S, \circ) = \text{Sym}(S)$ if and only if $(S, \circ)$ is either isomorphic or anti-isomorphic to one of the following types:*

- (i) *A right zero semigroup, i.e. $x \circ y = y$ for all $x, y \in S$.*
- (ii) *The idempotent quasigroup of order 3.*
- (iii) *The groupoid $(\{1, 2\}, \circ)$ where $x \circ 1 = 2$ and $x \circ 2 = 1$, for $x \in \{1, 2\}$.*

Furthermore, if $\text{Aut}(S, \circ)$ is triply transitive then $\text{Aut}(S, \circ) = \text{Sym}(S)$.

Groupoids with transitive automorphism group

We give some remarks on groupoids $(S, \circ)$ with transitive automorphism group that make computer searches for such groupoids more efficient.

Fix a groupoid $(S, \circ)$. For $s \in S$ define the sets

$$\begin{aligned} N_l(s) &:= \{x \in S \mid s \circ x = x\}, & N_r(s) &:= \{x \in S \mid x \circ s = x\}, \\ A_l(s) &:= \{x \in S \mid s \circ x = s\}, & A_r(s) &:= \{x \in S \mid x \circ s = s\}. \end{aligned}$$

These are the sets of all elements x for which s is left-neutral, right-neutral, left-absorbing and right-absorbing, respectively. Further let

$$\begin{aligned} n_l(s) &:= |N_l(s)|, & n_r(s) &:= |N_r(s)|, \\ a_l(s) &:= |A_l(s)|, & a_r(s) &:= |A_r(s)|. \end{aligned}$$

Lemma 3.8. *Let $(S, \circ)$ be a groupoid with transitive automorphism group. Then the cardinalities $n_l(s)$, $n_r(s)$, $a_l(s)$ and $a_r(s)$ are independent of s .*

Proof. Let $s, t \in S$, $s \neq t$. Then there exists $\varphi \in \text{Aut}(S, \circ)$ such that $\varphi(s) = t$. Now, for any $x \in S$ we have $s \circ x = x$ if and only if $\varphi(s) \circ \varphi(x) = \varphi(x)$, hence $x \in N_l(s)$ if and only if $\varphi(x) \in N_l(\varphi(s)) = N_l(t)$. This shows $\varphi(N_l(s)) = N_l(t)$ and thus $n_l(s) = n_l(t)$.

Similarly one observes $\varphi(N_r(s)) = N_r(t)$, $\varphi(A_l(s)) = A_l(t)$ and $\varphi(A_r(s)) = A_r(t)$, hence $n_r(s) = n_r(t)$, $a_l(s) = a_l(t)$ and $a_r(s) = a_r(t)$. $\square$

In the situation of Lemma 3.8 we may just write n_l , n_r , a_l and a_r for the corresponding cardinalities.

Lemma 3.9. *Let $(S, \circ)$ be a finite groupoid with transitive automorphism group. Then $n_l = a_r$ and $a_l = n_r$.*

Proof. For any groupoid $(S, \circ)$ it holds

$$\sum_{x \in S} n_l(x) = |\{(x, y) \in S^2 \mid x \circ y = y\}| = \sum_{y \in S} a_r(y) .$$

If $(S, \circ)$ has a transitive automorphism group it follows $|S|n_l = |S|a_r$ by Lemma 3.8. Hence, if $|S|$ is finite we have $n_l = a_r$. The proof of $a_l = n_r$ is very similar. $\square$

Corollary 3.10. *Let $(S, \circ)$ be a commutative finite groupoid with transitive automorphism group. Then $n_l = n_r = a_l = a_r$.*

Proof. We have $N_l = N_r$ and $A_l = A_r$ and thus $n_l = n_r$ and $a_l = a_r$. $\square$

Example 3.11. Let m be an odd positive integer and let $(\mathbb{Z}_m, +, \cdot)$ be the ring of integers modulo m . Define an operation $\circ$ on the set $\mathbb{Z}_m$ by

$$(a \circ b) := \frac{1}{2} \cdot (a + b) ,$$

where $\frac{1}{2}$ is the inverse of 2 in $\mathbb{Z}_m$.

Then $(\mathbb{Z}_m, \circ)$ is a commutative quasigroup with transitive automorphism group (and $n_l = n_r = a_l = a_r = 1$). In fact, for every $c \in \mathbb{Z}_m$ the translation map $\mathbb{Z}_m \rightarrow \mathbb{Z}_m$, $x \mapsto x + c$ is an automorphism of $(\mathbb{Z}_m, \circ)$. We get a generalised parasemifield $(\mathbb{Z}_m, \circ, +)$ where both operations are commutative, the addition is a quasigroup and the multiplication is a group.

4. Congruence-simple semirings

In this section congruence-simple semirings in which the multiplication is not necessarily associative will be studied.

Definition 4.1. A **semiring** is a ringoid $(S, +, *)$ in which the additive groupoid $(S, +)$ is associative and commutative.

Basic classification

Let $(S, +, *)$ be a semiring. For $a \in S$ and $n \in \mathbb{N} = \{1, 2, 3, \dots\}$ we define na recursively by $1a := a$ and $(n+1)a := na + a$.

Remark 4.2. Let $(S, +, *)$ be a semiring. For any $a, b \in S$ and $m, n \in \mathbb{N}$ we have $m(na) = (mn)a$ and $n(a + b) = na + nb$, which follows from the associativity and commutativity of the addition. Furthermore, it holds $a * (nb) = n(a * b)$ and $(na) * b = n(a * b)$, which is implied by the distributive laws.

Lemma 4.3. *Let $(S, +, *)$ be a semiring. Let the relation $\preceq$ on S be defined by*

$$a \preceq b \quad :\Leftrightarrow \quad \exists n \in \mathbb{N}, x \in S : nb = x + a$$

*and let $\rho := \preceq \cap \preceq^{-1}$. Then ρ is a congruence of the semiring $(S, +, *)$ and $\rho(a, 2a)$ for all $a \in S$.*

Proof. First we show that $\preceq$ is a preorder on S . Let $a, b, c \in S$. Since $2a = a + a$ we have $a \preceq a$. Now suppose $a \preceq b$ and $b \preceq c$. Then there exist $m, n \in \mathbb{N}$ and $x, y \in S$ such that $nb = x + a$ and $mc = y + b$. It follows $(nm)c = n(mc) = n(y + b) = ny + nb = ny + x + a$, so that $a \preceq c$.

Next we show that the preorder $\preceq$ is compatible with the semiring operations. Let $a, b \in S$ such that $a \preceq b$, and let $s \in S$. There exist $n \in \mathbb{N}$ and $x \in S$ such that $nb = x + a$. It follows $(n+1)(s+b) = ns + nb + s + b = ns + x + a + s + b = (ns + x + b) + s + a$, so that $s + a \preceq s + b$. Furthermore, $n(s * b) = s * (nb) = s * (x + a) = s * x + s * a$ and $n(b * s) = (nb) * s = (x + a) * s = x * s + a * s$, so that $s * a \preceq s * b$ and $a * s \preceq b * s$.

From what we have shown it follows that $\rho = \{(a, b) \in S^2 \mid a \preceq b \text{ and } b \preceq a\}$ is a congruence of the semiring $(S, +, *)$. Let $a \in S$. Then $a \preceq 2a$, since $1(2a) = a + a$, and $2a \preceq a$, since $3a = a + 2a$, hence $\rho(a, 2a)$. $\square$

Corollary 4.4. *Let $(S, +, *)$ be a congruence-simple semiring with an additively neutral element 0. Then the groupoid $(S, +)$ is either idempotent or a group.*

Proof. Consider the congruence ρ of Lemma 4.3. If $\rho = \text{id}_S$, then $2a = a$ for all $a \in S$, i.e. $(S, +)$ is idempotent. If $\rho \neq \text{id}_S$ then by congruence-simplicity it follows that $\rho = S \times S$, hence for all $a, b \in S$ there exists $n \in \mathbb{N}$ and $x \in S$ such that $nb = x + a$. In particular, for all $a \in S$ there exists $x \in S$ such that $0 = n0 = x + a$. Thus $(S, +)$ is a group. $\square$

Remark 4.5. Suppose that $(S, +, *)$ is a congruence-simple semiring without an additively neutral element such that $(S, +)$ is not idempotent. Along the same lines as the proof of Theorem 3.1 in [EHJK01] one can show that either the groupoid $(S, +)$ is cancellative or there is an absorbing element o of $(S, +)$ such that $2x = o$ for all $x \in S$.

Ideals and congruences

We examine the relationship between ideals, congruences and k -ideals (defined below). Afterwards we give a classification of k -ideal-simple semirings.

Lemma 4.6. *Let A be an ideal of a semiring $(S, +, *)$. Define a relation ρ_A on S by*

$$\rho_A := \{(x, y) \in S^2 \mid \exists a, b \in A : x + a = y + b\}.$$

*Then ρ_A is a congruence of the semiring $(S, +, *)$.*

Proof. The relation ρ_A is clearly reflexive and symmetric. Let $x, y, z \in S$ and suppose $\rho_A(x, y)$ and $\rho_A(y, z)$. Hence, there are $a, b, c, d \in A$ such that $x + a = y + b$ and $y + c = z + d$. It follows $x + a + c = y + b + c = y + c + b = z + d + b$. Since $a + c, b + d \in A$ we have $\rho_A(x, z)$ and thus ρ_A is an equivalence relation.

To show that ρ_A is a congruence, let $x, y \in S$ such that $\rho_A(x, y)$, and let $a, b \in A$ such that $x + a = y + b$. Further, let $z \in S$. Then $z + x + a = z + y + b$ and thus $\rho_A(z + x, z + y)$. Also, $z * x + z * a = z * y + z * b$ and $x * z + a * z = y * z + b * z$ with $z * a, z * b, a * z, b * z \in A$. Thus $\rho_A(z * x, z * y)$ and $\rho_A(x * z, y * z)$. $\square$

Definition 4.7. Let $(S, +, *)$ be a ringoid. A non-empty subset A of S is called a *k -ideal* of $(S, +, *)$ if A is an ideal of the ringoid $(S, +, *)$ such that $(A + A^c) \cup (A^c + A) \subseteq A^c$. Here, A^c denotes the complement $S \setminus A$ of A .

A ringoid $(S, +, *)$ is called *k -ideal-simple* if it has no proper k -ideals A with $|A| \geq 2$.

Lemma 4.8. *Let $(S, +, *)$ be a semiring with an additively neutral element 0 and let A be a k -ideal. Then the ρ_A -class $[0]_{\rho_A}$ of 0 equals A .*

Proof. We have $[0]_{\rho_A} = \{x \in S \mid \exists a \in A : x + a \in A\}$. This set equals A , since $a \in A$ and $x + a \in A$ implies $x \in A$ by the k -ideal property. $\square$

Corollary 4.9. *Let $(S, +, *)$ be a semiring such that $(S, +)$ has a neutral element. If the semiring is congruence-simple then it is k -ideal simple.*

Idempotent k -ideal-simple semirings

We consider now semirings with idempotent addition.

Remark 4.10. For any commutative idempotent semigroup $(S, +)$ there is an order relation $\leq$ on S defined by $\leq := \{(a, b) \in S^2 \mid a + b = b\}$. The partially ordered set $(S, \leq)$ is then a join-semilattice, where $a \vee b = a + b$ for all $a, b \in S$.

If $(S, +, *)$ is a semiring with idempotent addition and $\leq$ is defined as before, we have that $a \leq b$ implies $x * a \leq x * b$ and $a * x \leq b * x$ for any $a, b, x \in S$.

The following proposition gives an easily checkable criterion characterising finite additively idempotent k -ideal-simple semirings, which can be used to accelerate computer searches for congruence-simple semirings.

Proposition 4.11. *Let $(S, +, *)$ be a finite semiring with idempotent addition and let $\infty := \sum_{a \in S} a \in S$. Let M be the set of minimal elements in $(S, \leq)$. Then S is k -ideal simple if and only if for all $x \in S \setminus (M \cup \{\infty\})$ we have $\infty * x \not\leq x$ or $x * \infty \not\leq x$.*

Proof. First we show that a non-empty subset A of S satisfies

$$A + A \subseteq A \quad \text{and} \quad A + A^c \subseteq A^c$$

if and only if A is of the form $A_x := \downarrow x := \{a \in S \mid a \leq x\}$ for some $x \in S$.

Indeed, if $A + A \subseteq A$ and $A + A^c \subseteq A^c$ we let $x := \sum_{a \in A} a \in A$. We have $A = \downarrow x$, since each $a \in A$ satisfies $a \leq x$, and conversely, if $a \leq x$ then $a + x = x \in A$ and thus $a \in A$.

On the other hand, let $x \in S$ and $A = \downarrow x$. Then $A + A \subseteq A$, since $a, b \leq x$ implies $a \vee b \leq x$, for all $a, b \in S$. Furthermore, if $a \in A$ and $y \in S$ with $a + y \in A$ then $y \leq a + y \in A$, hence $y \in A$. Thus $A + A^c \subseteq A^c$.

Next for $x \in S$ we note that $S * A_x \subseteq A_x$ is equivalent to $\infty * x \leq x$, since $s * a \leq s * x \leq \infty * x$ for all $s \in S$ and $a \in A_x$. Similarly, $A_x * S \subseteq A_x$ if and only if $x * \infty \leq x$. Hence, A_x is a k -ideal if and only if $\infty * x \leq x$ and $x * \infty \leq x$.

We remark that $|A_x| = 1$ if and only if x is minimal in $(S, \leq)$, i.e. $x \in M$. In conclusion, there exists a proper k -ideal A with $|A| > 1$ if and only if there is $x \in S \setminus (M \cup \{\infty\})$ that satisfies the conditions $\infty * x \leq x$ and $x * \infty \leq x$. $\square$

Example 4.12. There are (up to isomorphism) five simple congruence-semirings $(\{0, 1, 2\}, +, *)$ of order 3 with idempotent addition and an *absorbing zero* 0, i.e. the element 0 is additively neutral and multiplicatively absorbing:

$+$	$\begin{array}{c ccc} & 0 & 1 & 2 \\ \hline 0 & 0 & 1 & 2 \\ 1 & 1 & 1 & 2 \\ 2 & 2 & 2 & 2 \end{array}$	$*$	$\begin{array}{c ccc} & 0 & 1 & 2 \\ \hline 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 2 & 0 & 2 & 2 \end{array}$	$*$	$\begin{array}{c ccc} & 0 & 1 & 2 \\ \hline 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \\ 2 & 0 & 2 & 2 \end{array}$
$*$	$\begin{array}{c ccc} & 0 & 1 & 2 \\ \hline 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 2 \\ 2 & 0 & 0 & 2 \end{array}$	$*$	$\begin{array}{c ccc} & 0 & 1 & 2 \\ \hline 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 2 \\ 2 & 0 & 1 & 2 \end{array}$	$*$	$\begin{array}{c ccc} & 0 & 1 & 2 \\ \hline 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 2 \\ 2 & 0 & 2 & 2 \end{array}$

The following table shows the number of congruence-simple semirings (up to isomorphism) of order n with idempotent addition and an absorbing zero:

n	2	3	4	5	6
# general	2	5	428	138 167	?
# commutative	2	1	21	715	59 640
# associative	2	0	0	0	1

The figures for associative semirings are taken from the classification [Zum08], the other numbers are results of a computer search program.

References

- [EHJK01] Robert ElBashir, Jan Hurt, Antonín Jančařík, and Tomáš Kepka, *Simple commutative semirings*, J. Algebra **236** (2001), no. 1, 277–306.
- [EK07] Robert ElBashir and Tomáš Kepka, *Congruence-simple semirings*, Semigroup Forum **75** (2007), no. 3, 589–609.
- [KK08] Vítězslav Kala and Tomáš Kepka, *A note on finitely generated ideal-simple commutative semirings*, Comment. Math. Univ. Carolin. **49** (2008), no. 1, 1–9.
- [KKK08] Vítězslav Kala, Tomáš Kepka, and Misolav Korbelař, *Notes on commutative parasemifields*, 2008, Preprint (10 pages), to appear in Comment. Math. Univ. Carolin.
- [Mon04] Chris Monico, *On finite congruence-simple semirings*, J. Algebra **271** (2004), no. 2, 846–854.
- [Tam63] Takayuki Tamura, *Some special groupoids*, Math. Japon. **8** (1963), 23–31.
- [Tam67] ———, *Notes on groupoids and their automorphism groups*, Proc. Japan Acad. **43** (1967), 843–846.
- [Zum08] Jens Zumbärgel, *Classification of finite congruence-simple semirings with zero*, J. Algebra Appl. **7** (2008), no. 3, 363–377.